


Copyright © 2022 National Council of State Boards of Nursing, Inc. (NCSBN*) All rights reserved. NCLEX*, NCLEX-RN* and NCLEX-PN* are registered All rights reserved. NCLEX*, NCLEX-RN* and NCLEX-PN* are registered trademarks of NCSBN* and may not be used or reproduced without written permission from NCSBN. No part of the these materials may be reproduced, stored in a retrieval system or transmitted in any form by any means (electronic, mechanical, photocopying, recording, or otherwise now known or to be invented) for any commercial or for profit use or purpose without written permission from NCSBN. Inquiries in writting to NCSBN Permissions, 111 E. Wacker Drive, Suite 2900, Chicago, IL 60601-4277 or via email at communications@ncsho.pre. communications@ncsbn.org. NCSBN


2




Why is the NCLEX changing? NCSBN


4


3





5 6






Measuring Clinical Judgment

Measuring Clinical Judgment

Topic The MOI Clinical Judgment Measurement Measurement Measurement
Moi Clinical Judgment Measurement
Moi Clinical Judgment


DOI PUBLICATION

Incsbn.org





9 10







11 12










15 16




Which of the following assessment findings require immediate follow-up? Select all that apply


| productive cough
| BP 9905.0 P 116, RR 24
| intermittent left shoulder pain
| ECG showing normal sinus thythm
| slightly diminished breath sounds on the left
| T 97 8° F (36 6° C), 0₂ saturation 99% on room air
| Hgb 9 gldL (90 g/L), HCT 27% (0.27), WBC 19,000imm³ (19.0 x 10³/L)
| tendemess upon palpation and duliness to percussion over the abdomen


| ncsbn.org | NCSBN | Locates appeared to the control of the cont

17 18











21 22





23 24





Take Action
Implementing the solution(s) that addresses the highest priorities.

Which intervention or combination of interventions is most appropriate?

How should the intervention(s) be accomplished (performed, requested, administered, communicated, taught, documented, etc.)?

Progress Notes

Client is post-op day #3 after a splenectomy and is able to ambulate in the corridor 3 to 4 times daily with minimal assistance. The client has clear breath sounds with a left chest tube in place attached to a closed-chest drainage system. Tidaling of the water chamber noted with deep inspiration. The client is refusing to use the incentive spirometer stating at causes left-sided chest pain. The client is refusing to use the incentive spirometer stating at causes left-sided chest pain. The client is utilizing prescribed patient-controlled analgesia (PCA). (device maximally every hour and continues to have intermittent nausea with some wormting. Adequate urine output. Abdominal surgical incision site with dressing is clean, dry, and intact with no erythema, edema or drainage noted to site.

elements from the scenario are what determines approach, Avoid memorized or "textbook" procedures. The item stem and/or the responses should include action verbs.

For "how" questions, ensure that specific


27 28







29 30





























39 40






41 42











45 46





47 48



